

## 1312 Somatosensory Therapists of Pain from 42 different countries

To **medical doctors**  
To **patients**

To **neuroscientists**  
To **therapists**

In 1992, the first communication about somatosensory rehabilitation of pain was done at the occasion of the 1<sup>st</sup> Congress of the swiss society for hand therapy. In 2001, this method was taught for the first time. On August 25<sup>th</sup> 2019, 1312 therapists, surgeons and medical doctors from **42 countries** have been trained to somatosensory rehabilitation of neuropathic pain.



● ≥ 300

● ≥ 100

● < 100

1	France	472
2	Canada (F)	249
3	Switzerland (F)	225
4	Switzerland (G)	139
5	The Netherlands	53
6	Belgium	32
7	Switzerland (I)	19
8	Canada (E)	18
9	India	17
10	Reunion Island	17
11	Germany	13
12	Luxemburg	8
13	Spain	8
14	Portugal	4
15	Greece	3
16	Finland	3

17	Turkey	3
18	Austria	3
19	Denmark	3
20	Roumania	2
21	Egypt	2
22	Italy	2
23	Israel	2
24	United-Kingdom	2
25	Czech Republic	1
26	Australia	1
27	Argentina	1
28	South Africa	1
29	USA	1
30	Brazil	1
31	Syria	1
32	Saudi Arabia	1

33	Japan	1
34	Estonia	1
35	Mauritius Island	1
36	Monaco	1
37	Martinique	1
38	Vietman	1
39	Tibet	1
40	Iran	1
41	China	1
42	Lebanon	1

**TOTAL**  
**1312**